

Exploring Microbial Dynamics in India's Rice Fields: Investigation of Plant Growth-Promoting Rhizobacteria for Sustainable Agriculture

Apoorv Pathak and Gurjeet Kaur*

Amity Institute of Biotechnology, Amity University Uttar Pradesh, Lucknow Campus 226028, India

*Corresponding author: gkaur@lko.amity.edu

Abstract

The interaction between plants and microorganisms in the rhizosphere plays a crucial role in soil health and plant development dynamics in agricultural environments. Plant Growth-Promoting Rhizobacteria (PGPR) have emerged as key players in enhancing plant growth, nutrient absorption, and stress tolerance, offering promising avenues for sustainable agriculture. This study investigates the microbial diversity in the rhizospheric soil of rice fields in India, focusing on identifying and characterizing putative PGPR strains.

To achieve this, rhizosphere soil samples were collected from rice fields, and specialized tools were used for extraction. The samples underwent controlled air-drying, sieving, and proper labeling before storage. PGPR isolation involved screening for symbiotic nitrogen-fixing microorganisms, non-symbiotic bacterial species, phosphate solubilizing microbes, and sulfate-reducing bacteria. Isolated PGPR strains were characterized using morphological, biochemical, and molecular analyses.

Molecular Evolutionary Genetics Analysis (MEGA) software and Basic Local Alignment Search Tool (BLAST) were used for evolutionary analysis and sequence similarity searches, respectively. The study revealed a significant microbial load in rhizospheric soil, indicating a thriving microbial community crucial for soil health and plant growth. Seasonal variations and agricultural practices were found to influence microbial abundance and composition.

Isolated PGPR strains exhibited diverse plant growth-promoting traits, highlighting their potential for sustainable agriculture. The study provides insights into the microbial dynamics in agricultural ecosystems and underscores the importance of PGPR in promoting soil health and plant productivity. By leveraging advanced analytical techniques and adopting sustainable agricultural practices, stakeholders can enhance agricultural sustainability and environmental stewardship.

Keywords: Plant Growth-Promoting Rhizobacteria (PGPR), Rhizospheric soil, Sustainable agriculture, Microbial diversity, Nitrogen fixation, Agricultural ecosystems.

Introduction

According to Bakker et al. (2013), complex interactions between plants and microorganisms influence soil health and plant development dynamics in agricultural environments, where the rhizosphere functions as a thriving microbial hotspot. Plant Growth-Promoting Rhizobacteria (PGPR), among the wide range of microorganisms that live in the rhizosphere, have become important for improving plant growth, nutrient

absorption, and stress tolerance (Kashyap et al., 2021). By encouraging environmentally friendly agricultural methods and lowering reliance on chemical pesticides and fertilisers, the investigation of PGPR communities offers great potential for sustainable agriculture. One of the most important staple crops in the world, rice (*Oryza sativa* L.) provides nourishment for a large proportion of the world's population. Rice farming is a fundamental agricultural activity in the agroecological environment of India, and it makes a significant contribution to both food security and economic stability. However, a number of issues, such as nutrient shortage, degraded soil, and environmental stresses, pose a threat to the sustainability and productivity of rice farming systems (Swarupa and Kiran, 2020). A vast reservoir of microbial variety may be found in the rhizospheric soil found in rice fields, and PGPR shows promise for transforming agricultural production and sustainability (Kour et al., 2019). Using the advantageous qualities of native PGPR strains presents a viable way to increase rice output, growth, and resistance to biotic and abiotic challenges. Important insights into the microbial dynamics underpinning plant-microbe interactions and their implications for agricultural sustainability can be gained by methodically screening and characterising PGPR strains from the rhizospheric soil of rice fields in the India region (Kumavat et al., 2022). The objective of this study project is to conduct a thorough investigation of the microbial diversity in the rhizospheric soil of India's rice fields, with an emphasis on identifying and describing putative PGPR strains. This study uses cutting-edge molecular and microbiological techniques to find native PGPR candidates with diverse traits that promote plant growth, such as nitrogen fixation, phosphate solubilization, phytohormone production, and phytopathogen-opposing biocontrol abilities.

Recent research has emphasised the importance of PGPR in improving crop resistance to diverse stresses, building on previous findings. For example, Kashyap et al. (2021) showed how PGPR helps rice plants absorb nutrients better and withstand stress, which increases agricultural output overall. In a similar vein, Kour et al. (2019) highlighted how PGPR may help lessen the negative impacts that environmental stresses have on agricultural ecosystems. Moreover, Swarupa and Kiran (2020) emphasised the significance of sustainable farming methods in tackling the issues of degraded soil and insufficient nutrients in rice production.

All things considered, incorporating PGPR-based techniques into India's rice farming systems has the potential to advance agricultural sustainability, boost crop yields, and lessen the negative effects of environmental stresses. By working together, academics, farmers, and policymakers may fully use PGPR to change agricultural practices in the direction of a more sustainable and environmentally friendly future.

Data Collection Methods

Sample collection from rice field rhizospheric soil is a meticulous process crucial for obtaining representative samples that reflect the diverse microbial communities and soil characteristics present in the field. The selection of sampling sites is conducted with utmost care, ensuring randomness to encompass the variability in soil texture, pH levels, and overall plant health across the field. This random selection approach guarantees a comprehensive understanding of the microbial dynamics within the rhizosphere.

To collect rhizosphere soil samples, specialized tools such as soil augers or corers are employed to extract soil cores from depths typically ranging from 5 to 10 cm. This depth range is chosen to capture the zone where root exudates and interactions between plant roots and soil microorganisms are most active, thereby

maximizing the representation of microbial populations. Each sample is carefully collected to avoid cross-contamination and ensure the integrity of subsequent analyses.

Following collection, the soil samples undergo controlled air-drying to prevent microbial activity and maintain sample stability. Air-drying is essential for preserving the microbial composition of the samples and preventing potential alterations in microbial populations due to environmental factors. Once dried, the samples are sieved through a 2 mm mesh to remove any debris or larger particles, ensuring homogeneity and consistency across samples.

Proper labeling of containers is essential to maintain sample traceability and prevent any confusion during further processing and analysis. Each container is labeled with relevant information, including sampling location, date, and any specific parameters of interest. Storage of the labeled containers at a temperature of 4°C helps preserve sample integrity and minimizes microbial activity until further analysis can be conducted.

The meticulous approach to sample collection outlined above ensures that the obtained rhizosphere soil samples are representative of the field's microbial diversity and soil characteristics. This comprehensive sampling strategy forms the foundation for subsequent analyses aimed at exploring the role of Plant Growth-Promoting Rhizobacteria (PGPR) and other microorganisms in rice cultivation and agricultural sustainability.

PGPR Isolation

The process of isolating and screening **Plant Growth-Promoting Rhizobacteria (PGPR)** involves a multifaceted approach aimed at identifying microbial strains with the potential to enhance plant growth, nutrient uptake, and stress tolerance. Among the various categories of PGPR, the screening focuses on symbiotic nitrogen-fixing microorganisms, non-symbiotic (free-living) bacterial species, phosphate solubilizing microbes, and sulfate-reducing bacteria. Each screening method targets specific microbial functions relevant to plant-microbe interactions and agricultural sustainability.

In the screening of symbiotic nitrogen-fixing microorganisms, such as *Rhizobium* and *Azotobacter*, rhizospheric soil samples are collected and processed. Serial dilutions of these samples are prepared, and aliquots are inoculated onto selective media such as YEMA Agar supplemented with Cycloheximide and Bromothymol Blue agar. This selective media allows for the identification of nitrogen-fixing microbes based on colony characteristics and their ability to produce acid or alkali. Nitrogen fixation is a crucial process in soil ecosystems, as it facilitates the conversion of atmospheric nitrogen into organic forms that are readily available to plants (Haerani et al., 2023).

In parallel, the screening of non-symbiotic (free-living) bacterial species from soil samples targets microbial strains capable of nitrogen fixation independently of plant roots. Non-rhizospheric soil samples are collected and processed similarly to rhizospheric samples, with serial dilutions and inoculation onto Ashby's Mannitol Agar medium. This screening method aims to identify free-living nitrogen-fixing microbes, which contribute to soil fertility and nutrient cycling, thus indirectly promoting plant growth and health (Sherpa et al., 2021; Bandeppa et al., 2019).

Additionally, the screening process includes the identification of phosphate solubilizing microbes, which play a crucial role in making phosphorus more accessible to plants. Non-rhizospheric soil samples are processed to prepare serial dilutions, and aliquots are inoculated onto Pikovskayas Agar medium. Phosphate solubilizing microbes are identified based on the visible appearance of zones of solubilization of inorganic phosphate around the colony. Phosphorus is an essential nutrient for plant growth and

development, and efficient phosphate solubilization by microbes can significantly enhance plant productivity (Gupta et al., 2022; Patel et al., 2022).

Furthermore, the screening process includes the identification of sulfate-reducing bacteria, which contribute to sulfur cycling and soil fertility. Non-rhizospheric soil samples are processed similarly to other screenings, with serial dilutions and inoculation onto Sulphate Reducing Agar medium. Sulfate-reducing microbes are identified based on the visible appearance of zones of solubilization of inorganic sulfate around the colony. Sulfur is an essential nutrient for plant metabolism and the synthesis of certain amino acids and proteins, highlighting the importance of sulfate-reducing bacteria in supporting plant growth (Lin et al., 2010; Ouattara and Jacq, 1992).

The screening process for PGPR encompasses a comprehensive exploration of microbial diversity within soil ecosystems, aiming to identify strains with multifaceted plant growth-promoting attributes. By isolating and characterizing these microbial strains, researchers can gain valuable insights into the intricate interactions between plants and microorganisms in agricultural ecosystems, paving the way for sustainable farming practices and enhanced crop productivity.

The comprehensive identification of Plant Growth-Promoting Rhizobacteria (PGPR)

The identification and characterization of **Plant Growth-Promoting Rhizobacteria (PGPR)** entail a comprehensive approach encompassing morphological, biochemical, and molecular analyses. Initially, isolated bacterial strains are cultured on selective agar media to assess their colony characteristics, including size, shape, color, margin, opacity, consistency, motility, and Gram characteristics (Hardiansyah, 2020; Naureen et al., 2005). These initial observations provide essential insights into the visual and structural features of the isolates, aiding in their preliminary classification and characterization.

Following the assessment of colony characteristics, the isolates undergo a series of biochemical assays to evaluate their metabolic capabilities. The **IMViC test**, comprising Indole, Methyl Red, Voges-Proskauer, and Citrate Utilization assays, is employed to detect specific enzymatic activities indicative of bacterial metabolic pathways (Das et al., 2019; Hamza et al., 2017; Mohite, 2013; Wagh et al., 2015). This battery of tests allows for the differentiation of bacterial species based on their metabolic profiles, providing crucial information about the physiological traits of the isolates.

Moreover, the isolates are subjected to the **carbohydrate fermentation test**, wherein they are inoculated into carbohydrate fermentation media containing various sugars. After incubation, fermentation patterns are observed to assess the strains' ability to utilize different carbohydrates as energy sources, aiding in their identification and classification (Deaker et al., 2008). Furthermore, the **catalase production assay** is employed to evaluate the isolates' ability to produce catalase enzyme, which neutralizes hydrogen peroxide. This assay involves adding hydrogen peroxide to bacterial suspensions and observing for immediate bubble formation, indicative of catalase activity (Reiner, 2010).

Additionally, the **amylase production assay** is conducted to detect the production of amylase enzyme by the isolates. They are inoculated onto starch agar plates and observed for the formation of hydrolysis zones after iodine staining. Amylase enzyme hydrolyzes starch into simpler sugars, reflecting the isolates' enzymatic capabilities and their potential roles in nutrient cycling and soil health (Visvanathan et al., 2020).

Beyond biochemical characterization, molecular techniques play a pivotal role in the identification and phylogenetic analysis of isolated PGPR strains. Specific isolates, such as *Azotobacter* spp., undergo molecular identification through nucleotide sequencing of the **16S rRNA gene**. This molecular-level

analysis provides insights into the genetic composition and evolutionary relationships of the isolates. Phylogenetic analysis elucidates the evolutionary history of the isolates by comparing their sequences with reference material from NCBI GenBank. Sequence alignment, phylogenetic tree construction, and assessment of tree reliability are conducted using software tools like Mega 11.0 (Aquilanti et al., 2020).




Figure 1 Schematic diagram represents the mechanism of PGPR (plant growth-promoting rhizobacteria).

Softwares which can be used in PGPR

The **Molecular Evolutionary Genetics Analysis (MEGA)** software stands as an indispensable tool for biologists engaged in the evolutionary analysis of DNA and protein sequence data. With its comprehensive suite of features, MEGA caters to a wide range of analytical needs, including sequence alignment, phylogenetic tree reconstruction, testing evolutionary hypotheses, estimating sequence divergences, accessing sequence data online, and generating detailed analysis descriptions (Kumar et al., 1994, 2008; Kumar and Dudley, 2007). Developed over the years, MEGA has become a cornerstone in molecular evolutionary analysis, offering maximum likelihood (ML) methods that aid in arranging DNA, RNA, and protein sequences to discern regions of similarity indicative of functional, structural, or evolutionary relationships.

The **Basic Local Alignment Search Tool (BLAST)** serves as another essential resource for biologists conducting sequence similarity searches. Accessible through a web interface or as a standalone tool, BLAST compares user queries to a vast database of sequences, identifying short matches between sequences and initiating alignments from these 'hot spots'. This powerful tool provides statistical information about alignments, such as the expect value or false-positive rate, enabling researchers to assess the significance of their findings (Gardy and Brinkman, 2006; Ye et al., 2006; Hamady et al., 2010). The National Center for Biotechnology Information (NCBI) hosts a BLAST server, where researchers can submit queries in FASTA format or sequence identifiers for comparison against the database to find the most similar sequences.

Through the integration of **MEGA** and **BLAST**, researchers gain access to robust analytical capabilities for exploring evolutionary relationships, identifying sequence similarities, and elucidating genetic and functional characteristics of biological sequences. These tools play pivotal roles in various fields of biological research, including evolutionary biology, genetics, molecular biology, and bioinformatics. By leveraging the power of computational tools like MEGA and BLAST, scientists can unravel the complexities of biological systems, uncover novel insights into evolutionary processes, and advance our understanding of the genetic basis of life.

Furthermore, the continuous development and refinement of these software platforms ensure that researchers have access to cutting-edge tools equipped with the latest algorithms and methodologies. As molecular and computational biology continue to evolve, MEGA and BLAST remain at the forefront, empowering scientists with the means to tackle increasingly complex biological questions and drive innovations in fields ranging from medicine to agriculture and beyond.

Characterization of Plant Growth Promoting Rhizobacteria (PGPR) isolates

In the realm of agricultural microbiology, the characterization of **Plant Growth-Promoting Rhizobacteria (PGPR)** holds immense significance, offering a pathway towards sustainable agriculture through the harnessing of beneficial microbial interactions. Within this domain, a meticulous assessment of various plant growth-promoting traits is paramount, providing insights into the functional capacities of microbial isolates and their potential applications in enhancing crop productivity and resilience. This comprehensive characterization often encompasses multiple assays targeting key traits such as indole acetic acid (IAA) production, nitrogen fixation, phosphate solubilization, and antimicrobial activity.

Indole Acetic Acid Production Assessment: Indole acetic acid (IAA), a phytohormone, plays a pivotal role in plant growth and development, influencing processes like cell elongation and root formation. The ability of PGPR isolates to produce IAA is evaluated using a colorimetric technique based on the Van Urk-Salkowski reagent and Salkowski's method (Ehmann, 1977). Cultured in yeast malt dextrose broth (YMD broth), isolates undergo incubation followed by centrifugation to collect the supernatant. Subsequent mixing with Salkowski's reagent allows for the quantification of IAA production through optical density (OD) readings at specific wavelengths using a spectrophotometer (Mohite, 2013).

Assessment of Nitrogen Fixation Activity by Deaminase Assay: Nitrogen is a vital nutrient for plant growth, and the ability of PGPR isolates to fix atmospheric nitrogen into biologically usable forms is a key trait contributing to their plant growth-promoting capabilities. To evaluate nitrogen fixation activity, isolates are inoculated onto Burk's medium, a nitrogen-free agar medium, under sterile conditions. Incubation at optimal temperatures allows for the observation of discernible growth, indicating proficient nitrogen fixation abilities in the isolates (Ha et al., 2018). Burk's medium provides a controlled environment conducive to accurate assessment and validation of nitrogen-fixing capabilities (Hartono et al., 2016).

Assessment of Phosphate Solubilization: Phosphorus is another essential nutrient for plant growth, often present in insoluble forms in soil. PGPR isolates capable of solubilizing phosphate compounds can enhance phosphorus availability to plants, thereby promoting growth and development. Assessment of phosphate solubilization involves the inoculation of isolates onto Pikovskayas Agar medium, followed by observation for the formation of clear zones around bacterial colonies, indicative of phosphate solubilization (Gupta et al., 2022; Patel et al., 2022).

Antimicrobial Activity Assessment: PGPR strains often exhibit antimicrobial properties, which can contribute to plant health by suppressing the growth of pathogenic microbes in the rhizosphere. The assessment of antimicrobial activity involves testing isolates against common plant pathogens using methods such as the agar well diffusion assay or the disc diffusion method. Zones of inhibition around bacterial colonies indicate the presence of antimicrobial compounds produced by the isolates (Santos et al., 2020; Tawfike et al., 2021).

Additional Traits and Characterization Techniques: In addition to the aforementioned traits, PGPR isolates can possess a myriad of other plant growth-promoting attributes, including siderophore production, ACC deaminase activity, and biofilm formation. Characterization techniques for these traits often involve specialized assays and biochemical analyses tailored to the specific functionalities of interest (Ma et al., 2020; Lopes et al., 2019).

Integration of Molecular Techniques: In recent years, the integration of molecular techniques has enhanced the characterization of PGPR isolates, allowing for a deeper understanding of their genetic composition and functional potential. Molecular methods such as PCR amplification of functional genes involved in plant-microbe interactions, metagenomic analysis of microbial communities, and transcriptomic profiling of gene expression patterns have provided valuable insights into the mechanisms underlying PGPR-mediated plant growth promotion (Bai et al., 2019; Niu et al., 2020).

Integration of Omics Approaches: Furthermore, omics approaches such as proteomics and metabolomics offer comprehensive insights into the biochemical pathways and metabolites associated with PGPR-mediated plant growth promotion. By elucidating the intricate networks of molecular interactions between PGPR and plants, omics-based studies contribute to a holistic understanding of the mechanisms driving plant-microbe interactions in agricultural ecosystems (Timmusk et al., 2011; Finkel et al., 2017).

Implications for Sustainable Agriculture: The comprehensive characterization of PGPR isolates and their plant growth-promoting traits has profound implications for sustainable agriculture. By leveraging the beneficial attributes of PGPR, farmers can reduce their reliance on chemical fertilizers and pesticides, mitigate environmental degradation, and enhance the resilience of crops to biotic and abiotic stressors. Moreover, the promotion of microbial diversity and ecosystem health through the application of PGPR-based interventions contributes to the long-term sustainability of agricultural systems (Bhattacharyya and Jha, 2012; Kumar et al., 2019). The characterization of PGPR isolates represents a critical step towards harnessing the potential of beneficial microorganisms for sustainable agriculture. Through a combination of traditional assays, molecular techniques, and omics approaches, researchers can gain a comprehensive understanding of the plant growth-promoting capabilities of PGPR strains and their implications for agricultural sustainability. By integrating these findings into agricultural practices, stakeholders can pave the way for a more resilient, productive, and environmentally friendly agricultural future.

In this comprehensive review article, we amalgamate various methodologies and findings to delve into the intricate dynamics of soil microbiota within agricultural ecosystems. Soil samples were initially subjected to serial dilution, reaching dilutions of up to 10^{-10} after being mixed with sterile distilled water. Aliquots from these dilutions were meticulously spread onto Petri dishes containing Plate Count Agar (PCA) and incubated for a duration ranging from 48 to 72 hours. The incubation period facilitated the growth of viable bacterial colonies, which were subsequently identified and counted, enabling the determination of colony-forming units (CFUs) per gram of soil (Rani and Reddi, 2012). These recorded

results were meticulously documented in Table 1 for further analysis, providing crucial insights into the microbial dynamics and ecological characteristics within the agricultural environment.

The assessment of microbial load revealed a significant range, spanning from 45 billion to 192 billion CFUs per gram of rhizospheric soil. Notably, the rhizospheric region exhibited remarkable density in terms of microbial load, suggesting a thriving microbial community inhabiting this niche. It is noteworthy that the microbial load may vary with seasonal changes, with the first month of summer potentially witnessing fluctuations in microbial abundance and diversity. This observation underscores the dynamic nature of soil microbial communities, influenced by various environmental factors and agricultural practices.

Furthermore, the analysis of microbial abundance and distribution within the rhizospheric soil provides valuable insights into the role of microorganisms in supporting plant growth, nutrient cycling, and overall soil health. The high microbial density observed in the rhizosphere underscores the significance of this microhabitat as a hotspot for microbial activity and interactions with plant roots. These interactions play pivotal roles in nutrient acquisition, disease suppression, and stress tolerance, ultimately contributing to enhanced plant growth and productivity in agricultural systems.

Moreover, understanding the seasonal variations in microbial abundance and composition is crucial for optimizing agricultural practices and promoting soil health and fertility throughout the year. Seasonal changes can influence microbial community dynamics, with fluctuations in temperature, moisture, and nutrient availability impacting microbial growth and activity. By elucidating these seasonal patterns, agricultural management strategies can be tailored to harness the beneficial effects of soil microorganisms and mitigate potential challenges associated with seasonal fluctuations.

In addition to seasonal dynamics, the microbial composition of the rhizospheric soil may also be influenced by agricultural practices such as crop rotation, tillage, and the application of fertilizers and pesticides. These practices can shape the diversity and functionality of soil microbial communities, with implications for soil fertility, plant health, and ecosystem resilience. Therefore, adopting sustainable agricultural practices that prioritize soil conservation and microbial biodiversity is essential for maintaining long-term agricultural productivity and environmental sustainability.




Figure 2 Soil Microbial Enzymes

Overall, the comprehensive analysis of soil microbial communities provides critical insights into the ecological functioning of agricultural ecosystems and underscores the importance of microbial diversity and activity in supporting soil health and plant productivity. By integrating advanced analytical techniques

and ecological principles, researchers can further elucidate the complex interactions between plants, microorganisms, and the soil environment, paving the way for innovative approaches to sustainable agriculture and environmental stewardship.

In the pursuit of isolating **non-symbiotic bacteria** capable of nitrogen fixation, a systematic screening process was employed. Initially, approximately 100 μL aliquots of 10^{-8} serially diluted soil samples were evenly spread onto Yeast Extract Mannitol Agar (YEMA Agar) plates. Subsequently, the Petri dishes were incubated for a period ranging between 48 to 72 hours under optimal conditions conducive to bacterial growth. Following incubation, well-isolated colonies were meticulously identified and recovered from the agar medium.

To further analyze the isolated colonies, a portion of the well-isolated colony was diluted in sterile saline water. A loopful suspension of the diluted culture was then streaked onto sterile Nutrient Agar plates using the streak plate method. This additional step facilitated the isolation of individual bacterial colonies and minimized the potential for mixed cultures, allowing for a more accurate assessment of colony morphology and characteristics.

The morphology and characteristics of the isolated colonies were meticulously observed and documented, as illustrated in Figures 2 and 3. Detailed observations included parameters such as colony size, shape, color, margin, opacity, and texture. These characteristics provide valuable insights into the diversity and phenotypic traits of the isolated bacterial strains, aiding in their classification and characterization.

This systematic screening approach aimed to identify and characterize non-symbiotic bacteria with potential **nitrogen-fixing capabilities**. Nitrogen fixation is a crucial process in agricultural ecosystems, as it facilitates the conversion of atmospheric nitrogen into biologically available forms that can be utilized by plants. By isolating and characterizing bacteria capable of nitrogen fixation, this study contributes to the understanding of microbial interactions and nutrient dynamics within agricultural ecosystems.

Furthermore, the identification of non-symbiotic bacteria with nitrogen-fixing capabilities holds significant implications for sustainable agriculture. These bacteria have the potential to enhance soil fertility, reduce the dependency on synthetic fertilizers, and promote environmentally friendly farming practices. By harnessing the natural ability of bacteria to fix nitrogen, farmers can improve crop yields, optimize nutrient management, and mitigate environmental pollution associated with nitrogen runoff.

Overall, the systematic screening of non-symbiotic bacteria for nitrogen fixation represents a critical step in advancing our understanding of microbial ecology and its applications in agriculture. By employing rigorous methodologies and detailed characterization techniques, researchers can uncover novel bacterial strains with beneficial traits, paving the way for innovative solutions to enhance agricultural sustainability and productivity.

In the pursuit of isolating **symbiotic bacteria** with potential nitrogen-fixing capabilities, a methodical screening process was employed. Initially, approximately 100 μL aliquots of 10^{-8} serially diluted soil samples were evenly spread onto Azotobacter Agar plates. These plates provided a selective medium conducive to the growth of symbiotic bacteria, particularly Azotobacter species. Following inoculation, the Petri dishes were incubated for a period ranging between 48 to 72 hours under optimal conditions to promote bacterial growth.

Following the incubation period, well-isolated colonies were meticulously identified and recovered from the agar medium. To further scrutinize the isolated colonies, a portion of the well-isolated colony was diluted in sterile saline water. Subsequently, a loopful suspension of the diluted culture was streaked onto

sterile Nutrient Agar plates using the streak plate method. This additional step allowed for the isolation of individual bacterial colonies and facilitated the observation of colony morphology and characteristics.

The morphological characteristics and features of the isolated colonies were meticulously observed and documented, as depicted in Figures 4 and 5. Detailed observations included parameters such as colony size, shape, color, margin, opacity, and texture. These observations provided valuable insights into the diversity and phenotypic traits of the isolated bacterial strains, aiding in their classification and characterization.

This methodical approach was employed to discern and characterize symbiotic bacteria with potential nitrogen-fixing capabilities. Symbiotic nitrogen fixation plays a crucial role in agricultural ecosystems by providing plants with a readily available source of nitrogen, thereby reducing the need for synthetic fertilizers and promoting sustainable farming practices. By isolating and characterizing symbiotic bacteria, this study contributes to the elucidation of microbial interactions and nutrient cycling dynamics within agricultural environments.

Furthermore, the identification of symbiotic bacteria capable of nitrogen fixation holds significant implications for agricultural sustainability. These bacteria form symbiotic relationships with plants, particularly legumes, and contribute to soil fertility by converting atmospheric nitrogen into a form that can be utilized by plants. By harnessing the natural ability of symbiotic bacteria to fix nitrogen, farmers can enhance crop yields, improve soil health, and reduce the environmental impact of agriculture.

Overall, the methodical screening of symbiotic bacteria for nitrogen fixation represents a critical step in advancing our understanding of microbial ecology and its applications in agriculture. By employing rigorous methodologies and detailed characterization techniques, researchers can uncover novel bacterial strains with beneficial traits, paving the way for innovative solutions to enhance agricultural sustainability and productivity.

In the quest to isolate **phosphate-solubilizing microbes**, a systematic screening approach was employed. Initially, approximately 100 μL aliquots of 10^{-8} serially diluted soil samples were evenly spread onto Pikovskaya's Agar Medium plates. Pikovskaya's Agar Medium is specifically designed to promote the growth of phosphate-solubilizing microorganisms by providing a suitable environment for the solubilization of insoluble phosphates. Following inoculation, the Petri dishes were incubated for a period of 6-7 days at 32°C , allowing for optimal bacterial growth and phosphate solubilization.

After the designated incubation period, well-isolated colonies were meticulously identified and recovered from the agar medium. The isolated colonies were subjected to detailed observation and documentation, as illustrated in Figures 6 and 7. Special attention was given to the zone of solubilization surrounding the colonies, which served as an indicator of phosphate-solubilizing activity. The size of the solubilization zone was recorded, typically measuring approximately 3-4 mm after an incubation period of approximately 7 days.

This systematic methodology aimed to isolate and characterize symbiotic bacteria with potential phosphate-solubilizing capabilities. Phosphorus is an essential nutrient for plant growth and development, but a significant portion of soil phosphorus exists in insoluble forms that are inaccessible to plants. Phosphate-solubilizing microbes play a crucial role in solubilizing these insoluble phosphates, making phosphorus more available to plants and contributing to improved nutrient uptake and plant growth. By isolating and characterizing phosphate-solubilizing microbes, this study contributes to the broader understanding of microbial interactions and nutrient cycling mechanisms within agricultural ecosystems. Phosphorus is one of the primary nutrients required for plant growth, and its availability in soil directly

impacts crop productivity and agricultural sustainability. Therefore, identifying microorganisms capable of solubilizing phosphorus can inform agricultural practices aimed at optimizing nutrient availability and enhancing crop yields while minimizing the need for external phosphorus inputs.




Figure 3 The main soil microbial enzymes and their roles in the soil. Acid/alkaline phosphatases (AP), arylsulphatase (ARS), β-1,4-glucosidase (BG), β(1-3) glucanase (β-Glu), catalases (CAT), cellobiohydrolase/exo- and endocellulases (CELB), dehydrogenases (DHA)

Furthermore, phosphate-solubilizing microbes can also play a role in soil fertility management and environmental sustainability. By promoting the solubilization of phosphorus, these microbes contribute to the efficient use of phosphorus resources and reduce the risk of phosphorus runoff and pollution in water bodies. Thus, the isolation and characterization of phosphate-solubilizing microbes have significant implications for sustainable agriculture and ecosystem health. Overall, the systematic screening of phosphate-solubilizing microbes represents a critical step in unraveling the complex interactions between microorganisms and nutrients in agricultural ecosystems. By employing rigorous methodologies and detailed characterization techniques, researchers can identify valuable microbial strains with the potential to enhance nutrient availability, soil fertility, and crop productivity, ultimately contributing to sustainable agriculture and food security.

In the pursuit of isolating **sulfate-reducing bacteria**, a systematic screening approach was employed. Initially, approximately 100 μL aliquots of 10⁻⁸ serially diluted soil samples were uniformly spread onto Sulphate Reducing Medium Agar plates. Sulphate Reducing Medium Agar provides an ideal environment for the growth of sulfate-reducing bacteria by supplying the necessary nutrients and sulfate source. Subsequently, the Petri dishes were subjected to an incubation period of 6-7 days at 32°C, facilitating optimal bacterial growth and sulfate reduction.

Following the designated incubation period, well-isolated colonies were meticulously observed and documented, as depicted in Figures 6 and 7. Special attention was given to the extent of solubilization surrounding the colonies, which served as an indicator of sulfate-reducing activity. The diameter of the solubilization zone was measured, typically yielding an average diameter of 3-4 mm after approximately 7 days of incubation.

This systematic approach aimed to isolate and characterize sulfate-reducing bacteria, which play a crucial role in the biogeochemical cycling of sulfur in soil ecosystems. Sulfur is an essential nutrient for plants, and its availability in soil is influenced by microbial processes such as sulfate reduction. Sulfate-reducing bacteria are capable of reducing sulfate to hydrogen sulfide, which can then be utilized by plants as a sulfur source. By isolating and characterizing sulfate-reducing bacteria, this study contributes to the comprehensive understanding of microbial diversity and biogeochemical processes within soil ecosystems.

Furthermore, sulfate-reducing bacteria are involved in various environmental processes, including sulfur cycling, metal bioremediation, and greenhouse gas production. These bacteria play a significant role in the sulfur cycle, influencing the transformation and availability of sulfur compounds in soil. Additionally, sulfate-reducing bacteria can contribute to the bioremediation of contaminated environments by reducing sulfate to sulfide, which can precipitate heavy metals and render them less toxic. Moreover, sulfate reduction is also associated with the production of hydrogen sulfide, a potent greenhouse gas, highlighting the importance of understanding the ecology and activity of sulfate-reducing bacteria in soil ecosystems. Overall, the systematic screening of sulfate-reducing bacteria represents a crucial step in unraveling the complex interactions between microorganisms and biogeochemical processes in soil ecosystems. By isolating and characterizing sulfate-reducing bacteria, researchers can gain insights into their ecological roles, contributions to nutrient cycling, and potential applications in environmental remediation and soil fertility management.

The comprehensive investigation of soil microbiota in agricultural ecosystems, particularly within the rhizospheric soil of rice fields in India, offers valuable insights into the intricate dynamics of microbial communities and their implications for agricultural sustainability. Through meticulous sampling, isolation, and characterization of microorganisms, this study sheds light on the role of Plant Growth-Promoting Rhizobacteria (PGPR) in enhancing soil health, nutrient cycling, and crop productivity.

Rhizospheric Microbial Dynamics: The assessment of microbial load in rhizospheric soil samples revealed a significant range, indicating the presence of a diverse and thriving microbial community. The high microbial density observed in the rhizosphere underscores its importance as a hotspot for microbial activity and interactions with plant roots. These interactions play pivotal roles in nutrient acquisition, disease suppression, and stress tolerance, ultimately contributing to enhanced plant growth and productivity in agricultural systems.

Seasonal Variations and Agricultural Practices: Seasonal changes and agricultural practices exert profound influences on soil microbial communities. The observed fluctuations in microbial abundance and composition underscore the dynamic nature of soil ecosystems, highlighting the need for adaptive agricultural management strategies. Understanding seasonal patterns and their impacts on microbial dynamics is essential for optimizing agricultural practices and promoting soil health throughout the year. Additionally, the adoption of sustainable agricultural practices, such as crop rotation and reduced tillage, can enhance soil fertility and microbial biodiversity, thereby supporting long-term agricultural productivity and environmental sustainability.

Isolation and Characterization of PGPR: The systematic screening and characterization of PGPR strains from rhizospheric soil samples involve multifaceted approaches aimed at identifying microbial strains with diverse plant growth-promoting attributes. Through selective media and biochemical assays, PGPR strains capable of nitrogen fixation, phosphate solubilization, and antimicrobial activity are

identified and characterized. Molecular techniques, including PCR amplification and sequence analysis, further elucidate the genetic composition and evolutionary relationships of isolated PGPR strains.

Implications for Sustainable Agriculture: The identification and characterization of PGPR strains hold immense implications for sustainable agriculture. By harnessing the beneficial attributes of PGPR, farmers can reduce reliance on chemical fertilizers and pesticides, mitigate environmental degradation, and enhance crop resilience to biotic and abiotic stressors. Furthermore, promoting microbial diversity and ecosystem health through PGPR-based interventions contributes to the long-term sustainability of agricultural systems.

Integration of Advanced Analytical Techniques: The integration of advanced analytical techniques, including molecular and omics approaches, enhances our understanding of the functional capacities and ecological significance of PGPR strains. Proteomics and metabolomics offer comprehensive insights into the biochemical pathways and metabolites associated with PGPR-mediated plant growth promotion, providing a holistic understanding of plant-microbe interactions in agricultural ecosystems.

Conclusion

In conclusion, the comprehensive investigation of soil microbiota and PGPR strains in agricultural ecosystems provides valuable insights into the ecological functioning of soil ecosystems and the potential for sustainable agriculture. By elucidating microbial dynamics, characterizing PGPR strains, and integrating advanced analytical techniques, researchers can pave the way for innovative approaches to agricultural sustainability and environmental stewardship. Through collaborative efforts among researchers, farmers, and policymakers, the harnessing of PGPR holds promise for transforming agricultural practices towards a more resilient, productive, and environmentally friendly future.

REFERENCES

1. Aquilanti, L., Favilli, F., & Clementi, F. (2004). Comparison of different strategies for isolation and preliminary identification of *Azotobacter* from soil samples. *Soil Biology and biochemistry*, 36(9), 1475-1483.
2. Bandeppa, S., Latha, P. C., Phule, A. S., Rajani, G., Prasad Babu, K. V., Barbadikar, K. M., Chandrakala, C., Prasad Babu,
3. M. B. B., Mandal, P. K., & Sundaram, R. M. (2019). Isolation, identification and characterization of efficient free-living nitrogen-fixing bacteria from rice rhizosphere ecosystem. *Journal of Rice Research*, 12(2), 39.
4. Bhardwaj, D., Ansari, M. W., & Sahoo, R. K. (2014). Tapping the rice rhizosphere for potent plant growth promoting bacteria. *FEMS microbiology ecology*, 88(1), 1-16.
5. Das, A., Kamal, S., Saha, S., & Sarkar, B. (2019). Effects of soil pH and temperature on the abundance of bacterial communities in the rhizosphere of rice. *Microbiology Open*, 8(4), 00700.
6. Das, S., Nurunnabi, T. R., Parveen, R., Mou, A. N., Islam, M. E., Islam, K. M. D., & Rahman, M. (2019). Isolation and characterization of indole acetic acid producing bacteria from rhizosphere soil and their effect on seed germination. *Int J Curr Microbiol Appl Sci*, 8(3), 1237-1245.
7. Deaker, R., Mijajlovic, G., & Casteriano, A. (2008). Estimating the most probable number of bacteria in multistrain biofertiliser inoculants using a multiple-tube fermentation test. Efficient nutrient use in rice production in Vietnam achieved using inoculant biofertilisers, 12-13.

8. Fatharani, R., & Rahayu, Y. S. (2018, November). Isolation and characterization of potassium-solubilizing bacteria from paddy rhizosphere (*Oryza sativa* L.). In *Journal of Physics: Conference Series* (Vol. 1108, No. 1, p. 012105). IOP Publishing.
9. Felsenstein, J. (1985). Phylogenies and the comparative method. *The American Naturalist*, 125(1), 1-15.
10. Gardy, J. L., & Brinkman, F. S. (2006). Methods for predicting bacterial protein subcellular localization. *Nature Reviews Microbiology*, 4(10), 741-751.
11. Glick, B. R. (2012). Plant growth-promoting bacteria: mechanisms and applications. *Scientifica*, 2012, 1-15.
12. Gupta, R., Kumari, A., Sharma, S., Alzahrani, O. M., Noureldeen, A., & Darwish, H. (2022). Identification, characterization and optimization of phosphate solubilizing rhizobacteria (PSRB) from rice rhizosphere. *Saudi Journal of Biological Sciences*, 29(1), 35-42.
13. Haerani, N., SYAM'UN, E. L. K. A. W. A. K. I. B., RASYID, B., & HARING, F. (2021). Isolation and characterization of N-fixing and IAA producing rhizobacteria from two rice field agro-ecosystems in South Sulawesi, Indonesia. *Biodiversitas Journal of Biological Diversity*, 22(5).
14. Hamady, M., Lozupone, C., & Knight, R. (2010). Fast UniFrac: facilitating high-throughput phylogenetic analyses of microbial communities including analysis of pyrosequencing and PhyloChip data. *The ISME journal*, 4(1), 17-27.
15. Hamza, T. A., Hussein, Z., Mitku, R., Ayalew, P., & Belayneh, T. (2017). Isolation and characterization of nitrogen fixing bacteria from rhizosphere soil collected from Shell Mele agricultural center, southern Ethiopia. *J AgricSci Food Technol*, 3(7), 117-124.
16. Hardiansyah, M. Y. H. M. Y. (2020). The Identification of Plant Growth Promoting Rhizobacteria from Thorny Bamboo Rhizosphere with 3% KOH Gram Test and Gram Staining Test. *International Journal of Applied Biology*, 4(2), 7-17.
17. Harrigan, W. F., & McCance, M. E. (2014). *Laboratory methods in microbiology*. Academic press.
18. Islam, S., Akanda, A. M., Prova, A., Islam, M. T., & Hossain, M. M. (2016). Isolation and characterization of plant growth-promoting rhizobacteria from cucumber rhizosphere and their effect on plant growth promotion and disease suppression. *Frontiers in microbiology*, 6, 1360.
19. Lin, H., Shi, J., Chen, X., Yang, J., Chen, Y., Zhao, Y., & Hu, T. (2010). Effects of lead upon the actions of sulfate-reducing bacteria in the rice rhizosphere. *Soil Biology and Biochemistry*, 42(7), 1038-1044.
20. Liu, L., Chen, S., Zhao, J., Zhou, X., Wang, B., Li, Y., ... & Huang, X. (2018). Watermelon planting is capable to restructure the soil microbiome that regulated by reductive soil disinfection. *Applied Soil Ecology*, 129, 52-60.
21. Majumdar, A., Dubey, P.K., Giri, B., Moulick, D., Srivastava, A.K., Roychowdhury, T., Bose, S. and Jaiswal, M.K., (2023). Combined effects of dry-wet irrigation, redox changes and microbial diversity on soil nutrient bioavailability in the rice field. *Soil and Tillage Research*, 232, p.105752.
22. Mohite, B. (2013). Isolation and characterization of indole acetic acid (IAA) producing bacteria from rhizospheric soil and its effect on plant growth. *Journal of soil science and plant nutrition*, 13(3), 638-649.
23. Naureen, Z., Yasmin, S., Hameed, S., Malik, K. A., & Hafeez, F. Y. (2005). Characterization and screening of bacteria from rhizosphere of maize grown in Indonesian and Pakistani soils. *Journal of Basic Microbiology: An International Journal on Biochemistry, Physiology, Genetics, Morphology, and Ecology of Microorganisms*, 45(6), 447-459.

24. Negi, R., Kaur, T., Devi, R., Kour, D., & Yadav, A. N. (2022). Assessment of nitrogen-fixing endophytic and mineral solubilizing rhizospheric bacteria as multifunctional microbial consortium for growth promotion of wheat and wild wheat relative *Aegilopskotschy*. *Heliyon*, 8(12).
25. Ouattara, A. S., & Jacq, V. A. (1992). Characterization of sulfate-reducing bacteria isolated from Senegal ricefields. *FEMS Microbiology letters*, 101(3), 217-228.
26. Patel, S., Prajapati, V., & Patel, P. (2022). Isolation and Screening of Mineral Phosphate Solubilizing Microorganisms. *Practical Handbook on Agricultural Microbiology*, 187-192.
27. Reiner, K. (2010). Catalase test protocol. *American society for microbiology*, 1-6.
28. Sherpa, M. T., Sharma, L., Bag, N., & Das, S. (2021). Isolation, characterization, and evaluation of native rhizobacterial consortia developed from the rhizosphere of rice grown in organic state Sikkim, India, and their effect on plant growth. *Frontiers in microbiology*, 12, 713660.
29. Singh, R. P., Jha, P. N., & Kumar, A. (2016). Microbial diversity in rice rhizosphere: a review. *African Journal of Microbiology Research*, 10(15), 534-546.
30. Sondo, M., Wonni, I., Klonowska, A., Kořta, K., & Moulin, L. (2023). Quantification of diversity sampling bias resulting from rice root bacterial isolation on popular and nitrogen-free culture media using 16S amplicon barcoding. *Plos one*, 18(4), e0279049.
31. Sood, Y., Singhmar, R., Singh, V., & Malik, D. K. (2023). Isolation and Characterization of Potential Potassium Solubilizing Bacteria with Various Plant Growth Promoting Traits. *Biosciences Biotechnology Research Asia*, 20(1), 79-84.
32. Tan, K.Z., Radziah, O., Halimi, M.S., Khairuddin, A.R., Habib, S.H., & Shamsuddin, Z.H. (2014). Isolation and Characterization of Rhizobia and Plant Growth-Promoting Rhizobacteria and Their Effects on Growth of Rice Seedlings. *American Journal of Agricultural and Biological Sciences*, 9(3), 342-360.
33. Visvanathan, R., Qader, M., Jayathilake, C., Jayawardana, B. C., Liyanage, R., & Sivakanesan, R. (2020). Critical review on conventional spectroscopic α -amylase activity detection methods: merits, demerits, and future prospects. *Journal of the Science of Food and Agriculture*, 100(7), 2836-2847.
34. Wagh, D. S., Shermale, R. N., & Mahure, B. V. (2015). Isolation and characterization of nitrogen fixing bacteria from agricultural rhizosphere. *IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS)* e-ISSN, 2319-2380.
35. Kashyap, A. S., Manzar, N., Rajawat, M. V. S., Kesharwani, A. K., Singh, R. P., Dubey, S. C., Pattanayak, D., Dhar, S., Lal,
36. S. K., & Singh, D. (2021). Screening and biocontrol potential of rhizobacteria native to gangetic plains and hilly regions to induce systemic resistance and promote plant growth in chilli against bacterial wilt disease. *Plants*, 10(10), 2125.
37. Swarupa, P., & Kiran, S. (2020). In vitro and in vivo characterization of plant growth promoting rhizobacteria isolated from rice rhizospheric soil. *International Journal of Botany Studies*, 5(5), 389-393
38. Kour, D., Rana, K. L., Yadav, N., Yadav, A. N., Kumar, A., Meena, V. S., ... & Saxena, A. K. (2019). Rhizospheric microbiomes: biodiversity, mechanisms of plant growth promotion, and biotechnological applications for sustainable agriculture. *Plant growth promoting rhizobacteria for agricultural sustainability: from theory to practices*, 19-65.

39. Kumawat, K. C., Nagpal, S., & Sharma, P. (2022). Potential of plant growth-promoting rhizobacteria-plant interactions in mitigating salt stress for sustainable agriculture: A review. *Pedosphere*, 32(2), 223-245.
40. Rani, M. U., Arundhathi, A., & Reddy, G. (2012). Screening of rhizobacteria containing plant growth promoting (PGPR) traits in rhizosphere soils and their role in enhancing growth of pigeon pea. *African Journal of Biotechnology*, 11(32), 8085- 8091.
41. Ha, T. T. T., Lam, T. T., Huyen, N. T., & Canh, N. X. (2018). Characterization and identification of nitrogenfixing bacteria isolated from agricultural soil. *Vietnam Journal of Science, Technology and Engineering*, 60(3), 48-54.
42. Hartono, H., Nurfitriani, N., Asnawati, F., Citra, H., Handayani, N. I., Junda, M., Ali, A., Hala, Y., & Jumadi, O. (2016). Ability of Ammonium Excretion, Indol Acetic Acid Production and Phosphate Solubilization of Nitrogen-fixing Bacteria Isolated from Crop Rhizosphere and Their Effect on Plant Growth. *ARPJ Journal of Engineering and Applied Sciences*, 11(19), 11735-11741.